Author Index

Åkesson, B., see Lindegård,B. 293
An, D., see Fang, X. 287
Anderson, R. A., see Brooks, C. J. W. 305
Ash, S., see Rustum, A. M. 209
Aubel, M. T.
——, Pesek, J. J. and Guiochon, G.
Characterization of an adamantyl-modified silica used as a stationary phase in high-performance liquid chromatography 11
Baeyens, W. R. G., see Lin Ling, B. 189
——, see Xianren, Q. 362
Baillet, A. E., see Taverna, M. 70
Balu, K., see Rustum, A. M. 209
Bauer, J. F.
——, Howard, S. and Schmidt, A.
High-performance liquid chromatographic determination of several quinolone antibacterials in medicated fish feed 348
Baylocq-Ferrier, D., see Taverna, M. 70
Brooks, C. J. W.
——, Cole, W. J. and Anderson, R. A.
Analytical gas chromatographic separations of diastereomeric *tert.*-butylmethoxyphenylsilyl ethers 305
Calabrò, G., see Micali, G. 317
Caldwell, G. W.
——, Masucci, J. A. and Jones, W. J.
Indirect thin-layer chromatography–fast atom bombardment and chemical ionization mass spectrometry determination of carbohydrates utilizing simple and rapid microtransfer techniques 377
Camps, F.
——, Coll, J., Pilar Marco, M. and Tomás, J.
Efficient determination of phytoecdysteroids from *Ajuga* species and *Polypodium vulgare* by high-performance liquid chromatography 199
Chou, T.-H., see Wu, S.-H. 325
Chui, S. H.
——, Lam, C. W. K., Lewis, W. H. P. and Lai, K. N.
High-performance liquid affinity chromatography for the purification of immunoglobulin A from human serum using jacalin 219
Cole, W. J., see Brooks, C. J. W. 305
Coll, J., see Camps, F. 199
Cserháti, T.
——, and Hauck, H. E.
Retention characteristics of CN, NH_2 and diol precoated high-performance thin-layer chromatographic plates in the adsorption and reversed-phase separation of some benzodiazepine derivatives 45
Currò, P., see Micali, G. 317
Dael, van M., see Smolders, E. 369
Danielson, N. D., see Wangsa, J. 171
Dannecker, W., see Schröder, B. 241
Dellacherie, J., see Görner, T. 309
Dewaele, C., see Lin Ling, B. 189
Do, L.
——, and Raulin, F.
Gas chromatography of the atmosphere of Titan. II. Analysis of low-molecular-weight hydrocarbons and nitriles with a potassium chloride-deactivated alumina PLOT capillary column 65
Dobarganes, M. C., see Márquez-Ruíz, G. 37
Doghieri, P., see Piretti, M. V. 334
Drevin, I.
——, Larsson, L., Eriksson, I. and Johansson, B.-L.
Chemical properties of and solute–support interactions with the gel filtration medium Superdex 75 prep grade 137
El Rassi, Z., see Nashabeh, W. 57
Ellingsen, T., see Kulin, L.-I. 1
Eriksson, I., see Drevin, I. 137
Fang, X.
——, Yu, B., Xiang, B. and An, D.
Application of pyrolysis–high-resolution gas chromatography–pattern recognition to the identification of the Chinese traditional medicine Mai Dong 287
Firmin, J. L.
High-performance liquid chromatographic analysis of octopinic acid, lysopine and nopalinic acid as sensitive indicators of *Agrobacterium*-induced crown gall tumours 343
Flodin, P., see Kulin, L.-I. 1
Fritsche, B., see Kennler, E. 383
Fuchiwaki, T., see Kawai, S. 29

Gaitonde, C. D.
—— and Pathak, P. V.
Separation and identification of impurities in the dye intermediate 8-amino-1-naphthol-3,6-disulphonic acid (H-acid) by high-performance liquid chromatography 330
—— and Pathak, P. V.
Capillary zone electrophoretic separation of chlorophenols in industrial waste water with on-column electrochemical detection 389
Galensa, R., see Krause, M. 147
Garrigues, P., see Wise, S. A. 111
Gill, R., see Smith, R. M. 97
Görner, T.
——, Dellacherie, J. and Perrut, M.
Comparison of helium head pressure carbon dioxide and pure carbon dioxide as mobile phases in supercritical fluid chromatography 309
Guiochon, G., see Aubel, M. T. 11
Hauck, H. E., see Cserháti, T. 45
Higashi, T., see Kawai, S. 29
Howard, S., see Bauer, J. F. 348
Hurtubise, R. J., see Mohseni, R. M. 19
Johansson, B.-L., see Drevin, I. 137
Jones, W. J., see Caldwell, G. W. 377
Jönsson, J. Å., see Lindegård, B. 293
Kawai, S., see Takayama, Y. 259
——, Fuchiwaki, T., Higashi, T. and Tomita, M.
High-performance liquid chromatographic determination of malonaldehyde using *p*-nitrophenylhydrazine as a derivatizing reagent 29
Kennler, E.
——, Schwer, Ch., Fritsche, B. and Pöhm, M.
Determination of arbutin in uvae-ursi folium (bearberry leaves) by capillary zone electrophoresis 383
Kihara, K., see Rokushika, S. 355
Knesel, L., see Šalamoun, J. 179
König, A., see Schoene, K. 279
Krause, M.
—— and Galensa, R.
Optical resolution of flavanones by high-performance liquid chromatography on various chiral stationary phases 147
Kulin, L.-I.
——, Flodin, P., Ellingsen, T. and Ugelstad, J.
Monosized polymer particles in size-exclusion chromatography, I. Toluene as solvent 1
Lai, K. N., see Chui, S. H. 219
Lai, S.-Y., see Wu, S.-H. 325
Lam, C. W. K., see Chui, S. H. 219
Lanuzzá, F., see Micali, G. 317
Lapouyade, R., see Wise, S. A. 111
Larsson, L., see Drevin, I. 137
Lazo, J. S., see Mistry, J. S. 86
Lecomte, L.
—— and Sinou, D.
Chiral sulphonated phosphines. IV[a]. High-performance liquid chromatographic separation of sulphonated phosphines 91
Leng, W.-X., see Rokushika, S. 355
Leoncini, R., see Terzuoli, L. 80
Lewis, W. H. P., see Chui, S. H. 219
Lin Ling, B.
——, Dewaele, C. and Baeyens, W. R. G.
Application of micro-scale liquid chromatography with fluorescence detection to the determination of thiols 189
Lin, S.-L., see Wu, S.-H. 325
Lindegård, B.
——, Mathiasson, L., Jönsson, J. Å. and Åkesson, B.
Controlled thermal degradation for identification and quantification of amine N-oxides in urine 293
Lou, X.
——, Sheng, Y. and Zhou, L.
Investigation of parameters in the separation of amino acid enantiomers by supercritical fluid chromatography 253
Macka, M., see Šalamoun, J. 179
Malmer, M. F.
——, and Schroeder, L. A.
Amino acid analysis by high-performance liquid chromatography with methanesulfonic acid hydrolysis and 9-fluorenylmethylchloroformate derivatization 227
Mangels, J., see Schröder, B. 241
Marinello, E., see Terzuoli, L. 80
Márquez-Ruíz, G.
——, Pérez-Camino, M. C. and Dobarganes, M. C.
Combination of adsorption and size-exclusion chromatography for the determination of fatty acid monomers, dimers and polymers 37
Masucci, J. A., see Caldwell, G. W. 377
Mathiasson, L., see Lindegård, B. 293
Matoušek, M., see Šalamoun, J. 179
Merckx, R., see Smolders, E. 371
Micali, G.
——, Lanuzzá, F., Currò, P. and Calabrò, G.
Separation of alkanes in *Citrus* essential oils by on-line coupled high-performance liquid chromatography–high-resolution gas chromatography 317
Mistry, J. S.
——, Sebti, S. M. and Lazo, J. S.
Separation of bleomycins and their deamido metabolites by high-performance cation-exchange chromatography 86

Mohseni, R. M.
—— and Hurtubise, R. J.
Reversed-phase chromatographic separation of selected hydroxyl aromatics with β-cyclodextrin as a mobile phase additive 19
Morikawa, M., see Takayama, Y. 259
Nashabeh, W.
—— and El Rassi, Z.
Capillary zone electrophoresis of pyridylamino derivatives of maltooligosaccharides 57
Neča, J.
—— and Vespalec, R.
Filled tubes as generation elements in electrokinetic detection in liquid chromatography 161
Nechvátal, M., see Šalamoun, J. 179
Ogura, T., see Yamagami, C. 123
Osselton, M. D., see Smith, R. M. 97
Pagani, R., see Terzuoli, L. 80
Pathak, P. V., see Gaitonde, C. D. 389, 330
Pérez-Camino, M. C., see Márquez-Ruíz, G. 37
Perrut, M., see Görner, T. 309
Pesek, J. J., see Aubel, M. T. 11
Pilar Marco, M., see Camps, F. 199
Piretti, M. V.
—— and Doghieri, P.
Separation of peracetylated flavanoid and flavanoid polyphenols by normal-phase high-performance liquid chromatography on a cyano-silica column and their determination 334
Pöhm, M., see Kennler, E. 383
Rassi, Z. El, see Nashabeh, W. 57
Raulin, F., see Do, L. 65
Rokushika, S.
——, Kihara, K., Subosa, P. F. and Leng, W.-X.
Ion chromatography of nitrite, bromide and nitrate ions in brine samples using a chloride-form anion-exchange resin column 355
Rustum, A. M.
——, Ash, S., Saxena, A. and Balu, K.
Reversed-phase high-performance liquid chromatographic method for the determination of soil-bound [^{14}C]atrazine and its radiolabeled metabolites in a soil metabolism study 209
Šalamoun, J.
——, Macka, M., Nechvátal, M., Matoušek, M. and Knesel, L.
Identification of products formed during UV irradiation of tamoxifen and their use for fluorescence detection in high-performance liquid chromatography 179
Sander, L. C., see Wise, S. A. 111
Saxena, A., see Rustum, A. M. 209
Schmidt, A., see Bauer, J. F. 346
Schoene, K.
——, Steinhanses, J. and König, A.
Sorption isotherms of organic vapours on Tenax 279
——, see Steinhanses, J. 273
Schröder, B.
——, Mangels, J., Selke, K., Wolpers, S. and Dannecker, W.
Comparative analysis of organic acids located on the surface of natural building stones by high-performance liquid, gas and ion chromatography 241
Schroeder, L. A., see Malmer, M. F. 227
Schwer, Ch., see Kennler, E. 383
Sebti, S. M., see Mistry, J. S. 86
Selke, K., see Schröder, B. 241
Sheng, Y., see Lou, X. 253
Sinou, D., see Lecomte, L. 91
Smith, R. M.
——, Westlake, J. P., Gill, R. and Osselton, M. D.
Retention reproducibility of basic drugs in high-performance liquid chromatography on a silica column with a methanol–high-pH buffer eluent. Effect of operating conditions on separations using an organic buffer 97
Smolders, E.
——, Van Dael, M. and Merckx, R.
Simultaneous determination of extractable sulphate and malate in plant extracts using ion chromatography 371
Steinhanses, J., see Schoene, K. 279
—— and Schoene, K.
Thermal desorption–gas chromatography of some organophosphates and S-Mustard after trapping on Tenax 273
Subosa, P. F., see Rokushika, S. 355
Takao, N., see Yamagami, C. 123
Takayama, Y.
——, Takeichi, T., Kawai, S. and Morikawa, M.
Behaviours of siloxane polymers containing phenyl or silarylene as stationary phases for high-temperature gas chromatography 259
Takeichi, T., see Takayama, Y. 259
Taverna, M.
——, Baillet, A. E. and Baylocq-Ferrier, D.
Identification of monosaccharides by high-performance liquid chromatography using methanolysis and a light-scattering detector 70
Terzuoli, L.
——, Pagani, R., Leoncini, R., Vannoni, D. and Marinello, E.
High-performance liquid chromatography of thiazolidinic compounds 80
Tomás, J., see Camps, F. 199
Tomita, M., see Kawai, S. 29
Ugelstad, J., see Kulin, L.-I. 1

Van Dael, M., see Smolders, E. 371
Vannoni, D., see Terzuoli, L. 80
Vespalec, R., see Neča, J. 161
Wangsa, J.
—— and Danielson, N. D.
Electrochemical detection for high-performance liquid chromatography using a Kel-F wax–graphite electrode 171
Westlake, J. P., see Smith, R. M. 97
Wise, S. A.
——, Sander, L. C., Lapouyade, R. and Garrigues, P.
Anomalous behaviour of selected methyl-substituted polycyclic aromatic hydrocarbons in reversed-phase liquid chromatography 111
Wolpers, S., see Schröder, B. 241
Wu, S.-H.
——, Lin, S.-L., Lai, S.-Y. and Chou, T.-H.
Separation of enantiomeric protected dipeptides by chiral high-performance liquid chromatography 325
Xiang, B., see Fang, X. 287
Xianren, Q.
—— and Baeyens, W.
Analysis of atmospheric precipitation by reversed-phase ion-pair chromatography 362
Yamagami, C.
——, Ogura, T. and Takao, N.
Hydrophobicity parameters determined by reversed-phase liquid chromatography. I. Relationship between capacity factors and octanol–water partition coefficients for monosubstituted pyrazines and the related pyridines 123
Yu, B., see Fang, X. 287
Zhou, L., see Lou, X. 253